

# On the use of cerebral and emotional measurements for the evaluation of workload during car driving

Enrica Modica<sup>\*a</sup>, Dario Rossi<sup>\*a</sup>, Patrizia Cherubino<sup>a,b</sup>, Arianna Trettel<sup>b</sup>, Wanzeng Kong<sup>c, a</sup>, G. Borghini<sup>a,b</sup>, F. Babiloni<sup>a,b</sup>, P. Aricò<sup>a,b</sup>, Alfredo Colosimo<sup>d</sup>, Giovanni Vecchiato<sup>e</sup> and Anton Giulio Maglione<sup>a,b</sup>

<sup>a</sup>Dept. Molecular Medicine, University Sapienza, Rome, Italy

<sup>b</sup>BrainSigns srl, Rome, Italy

<sup>c</sup>Dept. Computer Science, Hangzhou Dianzi University, Hangzhou, China

<sup>d</sup>Dept. of Anatomical, Histological, Forensic & Orthopedic Sciences, University Sapienza, Rome, Italy

<sup>e</sup>CNR Institute of Neuroscience – URT, Parma, Italy

Correspondence: F. Babiloni, Dept. Molecular Medicine, University Sapienza, viale Regina Elena 291, 00161, Rome, Italy. E-mail: fabio.babiloni@uniroma1.it, phone +39-3287697914

**Abstract.** It has been reported as the measurements of cerebral activity by EEG and other neurophysiologic variables such as EKG and eye blinks could be extremely useful in the characterization of the “internal” state of pilots and air traffic controllers. Here, we report the results from a study aimed to use a set of validated neurometric indicators for the study of the mental workload that could occur during simulated car drive in a group of healthy subjects. In addition, it was also investigated the insurgence of sleepiness during such driving activity. Results obtained suggest that the workload index is sensitive to the mental efforts of the driver during the different drive tasks performed. In addition, results suggested as HR increases during the execution of the difficult driving tasks while instead it decreases at the insurgence of the drowsiness. The proposed workload index could be then used in a near future to assess on-line the mental state of the driver during a drive task.

**Keywords:** EEG, PSD, Workload, Heart Rate, Eye Blinks rate

## 1. Introduction

Health Organization (WHO) indicates that road accidents are the ninth cause of death among young people aged between 18 and 29 years. Often this death occurs from a lapse of attention of the driver due to the onset of mental fatigue or induced by high cerebral workload. So the workload, fatigue and sleepiness have an important role in road safety. In fact, mental fatigue and drowsiness leading to a lowering of the level of attention and appeared to be among the main causes of road accidents. Several studies have shown a relationship between the cognitive status of the subject and his/her EEG activity during simple sensory -motor or cognitive tasks [Berka et al., 2007; Borghini et al., 2011; Fairclough and Tattersal, 2005; Lal and Craig, 2000; Baulk et al., 2001]. In particular, the EEG was found to be sensitive to changes in alertness and able to predict a decline in performance due to the increase in mental workload [Berka et al., 2007; Borghini et al., 2011; Fairclough and Tattersal, 2005; Lal and Craig, 2000; Baulk et al., 2001; Bach et al., 2009]. In these studies, it was showed an increase of the EEG power spectral density PSD in the theta frequency band over the prefrontal, frontal and parietal cortical areas, during difficult drive [Simon et al., 2011; Papadelis et al., 2007]. These changes have been often associated to a simultaneous reduction of the EEG PSD in alpha band over the parietal cortical areas [Papadelis et al., 2007; Klimesch, 1999; Borghini et al., 2012].

From these results provided by the literature it can be hypothesized that the generation of an efficient workload index for the driving task could be based on the simultaneous occurrence of the above mentioned changes in EEG PSD as well as with the use of the other autonomic measurement variables. In fact, other lines of experimental evidences suggested as the heart rate (HR) and the eye

blink duration (EBR) measurements could be correlated with the variation of the cognitive efforts during the drive task. In particular it is shown that an increase of HR could be indicative of an increased workload while the eye blink is inversely correlated with the changes in mental load of the drivers [Lal and Craig, 2000; Baulk et al., 2001; Bach et al., 2009; Simon et al., 2011; Papadelis et al., 2007; Klimesch, 1999; Borghini et al., 2012]. Thus, it can be hypothesized that EEG, EBR and HR could be employed in the generation of a workload index that will track the perceived difficulty of the driving task from the drivers. To check such hypothesis, we generated an experimental paradigm in which simulated driving tasks were proposed to a group of normal subjects. Such driving tasks have different level of difficulty or monotony, and the measure of EEG, HR and EBR variables were concurrently performed. The observed results for this experiment reported in the following, suggested that it is possible derive an efficient workload index on the base of such cerebral and autonomic measurements for the drive

## **2. Material and Methods**

### **2.1. Subjects**

The experimental group was composed by 20 healthy subjects (9 female, 11 male), all within 24 and 30 years old. All of them usually drive a car with the manual gear, of the same type of the gear proposed during the drive simulations. To all the subjects were prohibited to drink alcohol and to have heavy meals for one day prior to the measurements. Subjects were also asked to avoid caffeine up to 5 hours before the experiment. Each subjects gave his/her explicit informed consent to the experimental procedure.

### **2.2. Experimental protocol**

The experiments were performed between 2 PM and 5 PM. In this part of the day time sleepiness tends to increase [Baulk et al., 2001]. The experimental protocol is developed along 1 day. The simulated drive game used is Need For Speed and the Spa – Francorchamps (Belgium) track and the car by which performing the driving tasks was the Alfa Romeo – Giulietta QV (1750 TBi, 4 cylinders, 235 HP) have been chosen. The night condition for the drive task was also selected. In order to modulate the difficulty of the proposed driving task, a simultaneous attentional and vigilance task (TAV) was added concurrently to the drive task. This TAV task works as follows: an alert stimuli, a white “X”, was presented on a monitor placed about 1 m from the subject, avoiding the interference with the main screen where the game visualized. Subjects are asked to press the button #1 when the X appears on the screen, during the drive. The audio vigilance stimuli were generated by two speakers placed on the left and on the right side of the driver. A succession of frequent (with a 95% probability rate) and rare (5% probability) tones at different acoustic frequency were continuously delivered to the subjects. They are asked to press the button #2 when the rare stimulus occurs. These tasks (TAV) could be generated at different levels of difficulty, depending on the overall frequency of the visual and acoustic stimulations. The alternating tone sequence of the vigilance task could simulate the car’s radio or engine noise while the target tone sequence could be an unexpected sound (e.g. phone calling). The alert’s stimuli simulated instead the traffic jam, for example, the traffic lights, the pedestrians, other cars or other uncontrollable traffic agents [Bach et al., 2009]. After training of half hour the subject had to drive the simulated cars in two conditions: the first by moving the car in the selected circuit at a predetermined speed. Such condition were named “warm up” (WUP) and serve to collect the baseline for the cerebral and autonomic signals. The second drive condition, named “performance” (PERFO), required to the driver to race at a speed higher than that held on the warm up period. On this driving task it have been superimposed the occurrence of the different TAV tasks. In particular, five TAV tasks with increasing difficulty levels due to the increase of frequency of stimuli were employed during the experiments. Such TAV tasks are presented in a pseudo-random order to the subjects such as: TAV3, TAV1, TAV5, TAV2, TAV4, where TAV1 is the easier task and the TAV5 the most difficult one. Finally, the last condition in which the EEG and autonomic parameters of the subjects were collected was instead a monotonous night driving, in which the subject had to drive, on the same track without exceeding the speed of 70 (Km/h). With the last task we wanted to induce hypovigilance or drowsiness, that effectively occurred in all the subjects during the experiments. At the end of each task of the experiment subjects were asked to fill out the NASA-TLX questionnaire, that provides a standard measurement of

the perceived mental load during the task by the subjects. Errors performed by the subjects during the performance of the different TAV tasks during the simulated drive were also analyzed off-line.

### **2.3. EEG and Physiological recording**

A digital ambulatory monitoring system (Brain Products GmbH, Germany) are used to record EEG and physiological signals. The signals are acquired with a sampling frequency of 200 Hz and are collected simultaneously during the experiment. To remove power interference was applied a 50-Hz notch filter. The earlobes are used as reference and the impedances were maintained around 10 (k $\Omega$ ). The EEG recording was filtered with a band pass filter (1-30 Hz) and then Independent Component Analysis (ICA) was used to remove the artifacts and separate blink component from the data. The blink component are used for the estimation the eye blinks rate (EBR). The EKG raw signal was filtered with pass-band filter (low-pass filter cut-off frequency: 10 Hz, high-pass filter cut-off frequency: 0.5 Hz) used for estimate the heart rate (HR).

### **2.4. Neurophysiologic indexes for characterizing drowsiness and mental fatigue.**

It has been suggested as the onset of the state of drowsiness could be indexed by the appearance on the EEG traces of the alpha spindles [Simon et al., 2011]. Such spindles are defined as short bursts in the alpha band of EEG. and consist of 12-14 Hz waves that occur for at least 0.5 (s). It has been also observed as those EEG alpha bursts are also correlated with driving errors. We controlled the occurrences that these EEG events before the driving task in which the drowsiness was induced. In this work, the workload index (IWL) has been defined as the ratio between the EEG PSD in theta band over the central frontal area (Fz) and the EEG PSD in alpha band over the central parietal area (Pz). The EEG bands have been defined by using the Individual Alpha Frequency (IAF) [Klimesch, 1999; Borghini et al., 2012; Millan et al., 2000; Zavaglia et al., 2006; Babiloni et al., 2002; Babiloni et al., 2004; Babiloni et al., 2001]. The IWL, HR and EBR values have been subjected to the Analysis of Variance (ANOVA) in order to investigate changes of their values across the modulation of the TAV tasks. Factors for the ANOVA are TASK (with 6 levels) and GENDER (male, female). For HR and EBR indexes are used Z-score value, instead for IWL index are used r-squared values.

## **3. Results**

### **3.1. Workload index**

The group values of the estimated cerebral IWL across the different driving conditions is presented in Fig.2, in which the gender of the investigated sample it is also taken into account. The EEG data are calculated with r-squared. R-squared is a statistical measure computed over a pair of sample distributions, giving a measure of how strongly the means of the two distributions differ in relation to variance.




Figure 1. The cerebral IWL for men (blue line) and for women (red line) across the different drivin tasks proposed. For the all population investigated, as the difficult of tasks increases the cerebral IWL increases while it decreases during the drowsy condition (DROW).

Fig.2 shows an example of how the employed workload index changed during the different experimental conditions. Woman and men show the same trend across the different tasks. The TAV tasks are statistically different from the DROW condition ( $F=4.54$ ,  $p=0.004$ ). Application of the post-hoc Duncan tests revealed that the IWL average values for the population investigated discriminate statistically the difficult task (TAV5, TAV4, TAV3) from the easier tasks (TAV1, TAV2), in the female sample population. In the male such trend doesn't reach the statistical significance ( $p=0.1$ ).

### 3.2. Heart and eye blinks rate

The eye blinks rate (EBR) variable shows a statistical significant difference on average between the TAV conditions and the DROW one ( $p<0.05$ ). In fact, Fig.2 shows the EBR average values across the different TAV conditions and the DROW one. Male sample presented EBR values higher in DROW conditions than in the other ( $p<0.001$ ).




Figure 2. The EBR for men (blue line) and for women (red line) has been demonstrated to be different, especially in the drowsy condition (DROW). The EBR for men is higher in the drowsy condition and smaller in the TAV conditions.

The hearth rate (HR) variable showed a statistical difference between TAV and DROW condition ( $p < 0.05$ ), as well as for the gender factor (Fig.3).




Figure 3. HR average values for the male and female populations across the different TAV and the DROW condition.

In fact, in male population the HR values are higher than in the female population.

### 3.3. NASA TLX

The result obtained from NASA TLX test represented in Fig. 4 show that the subjective workload perceived in the analyzed population correlate with the level of the drive difficulty. In fact, such increase occur during the TAV tasks that are more difficult (TAV5,TAV3,TAV4) while it is less prominent during the easier tasks (TAV1,TAV2). Both populations (male, female) show statistical difference between TAV and DROW condition ( $p < 0.02$ ).




Figure 4. This picture shows the subjective workload evaluation as performed by filling the NASA-TLX questionnaire. The blue line represents NASA TLX results for men and the red line the NASA-TLX for women. The higher the NASA-TLX index the higher the perceived workload. Both women and men feel higher values of workload during the TAV conditions and smaller during the drowsy condition (DROW).

## 4. Discussion

In this paper it was showed that different cerebral (IWL) and autonomic (HR, EBR) indexes modulate with the levels of difficulty for a series of simulated drive tasks, at the group level. In addition, it is also observed as such results appear to be dependent to the gender of the relatively young population investigated. A common trend for IWL, HR and EBR indexes has been found; the higher task's difficulty, the higher the IWL and HR values, the NASA-TLX scores, and the driving errors (off road). In addition, the EBR decreased with the increase of difficulty and it decreased during the monotonous condition. However, such differences have been observed at the group level. The female group investigated show less amplitude in each indices considered.

This study suggests that EEG data can be used in the development of an on-line cerebral mental workload index (IWL). It also suggests that the integration of IWL information with those deriving from the autonomic indexes, such as heart rate and the eye blinks rate, could be important. The idea that the workload index could be estimated in a near future in "real-time" by a mixed system that includes EEG, HR and EBR signals inside the car appears feasible in the next few years. Such system will return feedback to the driver about his/her internal cognitive conditions during the execution of the drive task.

## Acknowledgements

This work was partly supported by Major International Cooperation Project of Zhejiang Province (Grant No. 2011C14017). It is also supported by BrainSigns srl, with a grant from Regione Lazio, FILAS, named "BrainTrained", CUP F87I12002500007 and by the Italian Minister of Foreign Affairs with a bilateral project between Italy and China "Neuropredictor".

## References

- Babiloni C., Babiloni F., Carducci F., Cappa S., Cincotti F., Del Percio C., "Human cortical EEG rhythms during long-term episodic memory task. A high-resolution EEG study of the HERA model", *Neuroimage* 21 (4), 1576-1584, 2004.
- Babiloni C., Babiloni F., Carducci F., Cincotti F., Rosciarelli F., "Human brain oscillatory activity phase-locked to painful electrical stimulations: A multi-channel EEG study", *Human brain mapping* 15 (2), 112-123, 2002.
- Babiloni C., Babiloni F., Carducci F., Cincotti F., Rosciarelli F., Arendt-Nielsen L., Chen A., Rossini PM., "Mapping of early and late human somatosensory evoked brain potentials to phasic galvanic painful stimulation", *Human brain mapping* 12 (3), 168-179, 2001.
- Bach K. M., Jæger M. G., Skov M. B., Thomassen N. G., "Interacting with In - Vehicle System: Understanding, Measuring, and Evaluating Attention", *HCI 2009 - People and Computers XXIII - Celebrating people and technology*, pp. 453 - 462.
- Baulk S. D., Reyner L. A., Horne J. A., "Driver Sleepiness - Evaluation of Reaction Time Measurement as a Secondary Task", *Sleep*, vol. 24, no. 6, pp. 695-698, 2001.
- Berka C., Levendowsky D. J., Lumicao M. N., Yau A., Davis G., Zivkovic V. T., Olmstead R. E., Tremoulet P. D., Craven P. L., "EEG correlates of task engagement and mental workload in vigilance, learning, and memory tasks", *Aviat Space Environ Med*, vol. 78, no. 5, pp. B231 - B244, 2007.
- Borghini G, Astolfi L, Vecchiato G, Mattia D, Babiloni F. "Measuring neurophysiological signals in aircraft pilots and car drivers for the assessment of mental workload, fatigue and drowsiness" *Neurosci Biobehav Rev*. 2012 Oct 30.
- Borghini G., Isabella R., Vecchiato G., Toppi J., Astolfi L., Caltagirone C., Babiloni F., "Brainshield: HREEG study of perceived pilot mental workload", *Italian journal of aerospace medicine*, vol. 5, pp. 34 - 47, 2011.
- Fairclough S. H., Tattersall L., "The influence of task demand and learning on the psychophysiological response", *International Journal of Psychophysiology*, vol. 56, no. 2, pp. 171 - 184, 2005.
- Klimesch W., "EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis", *Brain Research Reviews*, vol. 29, pp. 169-195, 1999.
- Lal S. K. L., Craig A., "Driver Fatigue: psychophysiological effects", *The Fourth International Conference on Fatigue and Transportation*, Australia, 2000.
- Millan J, Mourino J., Babiloni F., Cincotti F., Varsta M., Heikkonen J. "Local neural classifier for EEG-based recognition of mental tasks *Neural Networks*", 2000. *IJCNN 2000, Proceedings of the IEEE-INNS-ENNS International Joint Conference*, 2000.

Papadelis C., Chen Z., Kourtidou-Papadeli C., Bamidis P. D., Chouvarda I., Bekiaris E., Maglaveras N., “*Monitoring sleepiness with on – board electrophysiological recordings for preventing sleep - deprived traffic accidents*”, *Clinical Neurophysiology*, vol. 118, pp. 1906 – 1922, 2007.

Simon M., Schmidt E. A., Kincses W. E., Fritzsche M., Bruns A., Aufmuth C., Bogdan C., Rosenstiel W., Schrauf M., “*EEG alpha spindle measures as indicators of driver fatigue under real traffic conditions*”, *Clinical Neurophysiology*, vol. 122, no. 6, pp. 1168 – 1178, 2011.

Zavaglia M, Astolfi L., Babiloni F., Ursino M., “*A neural mass model for the simulation of cortical activity estimated from high resolution EEG during cognitive or motor tasks*”, *Journal of neuroscience methods* 157 (2), 317-329, 2006.